

Acceptable proof of immunity

FOR MEASLES

(Rubeola)

2 Doses of live measles vaccine, the first one administered on or after the first birthday, and the second after 15 months of age. There must be an interval of at least 30 days between the first and second doses of vaccine.

OR

Serological proof of immunity (Blood test for presence of measles antibody specify Rubeola).

OR

physician statement of prior disease history.

FOR RUBELLA

German Measles

(3 day
measles)

1 dose of vaccine given on or after the first birthday.

OR

Serological evidence of immunity (blood test for presence of rubella antibodies).

A physician statement of disease history is NOT sufficient for proof of immunity against rubella.

FOR MUMPS

1 dose of vaccine on or after the first birthday is required.

OR

Serological proof

OR

physician statement of prior disease history.

Month/day/year of immunizations required! Divisions of Public Health will not accept month/year without the date of month recorded.

IDEAS FOR OBTAINING IMMUNIZATION DATA

1. High School Health Records
2. Division of Public Health
3. Armed Services Health Records
4. Migrant Health Records (Shot Records)
5. Union Health Record
6. Signed Immunization Transfer Card
7. Immunization Portion of a Passport
8. Immunization Record Card
9. Pediatrician

If the student cannot provide any acceptable forms of proof, and if there are not religious or medical contraindications, re-immunization is required. However, serologic evidence of immunity is another option.